arXiv:0910.2870v2 [cond-mat.stat-mech] 17 May 2010

An extension of the equilibrium fluctuation relations

L. Velazquez[1,2] **and S. Curilef**[2]

Abstract. Recently, we have derived a generalization of the known canonical fluctuation relation between the heat capacity and the energy fluctuations $C = \beta^2 \left\langle \delta U^2 \right\rangle$, which is able to describe the existence of macrostates with negative heat capacities $C < 0$. In this work, we extend our previous analysis to the generalization of some other equilibrium fluctuation relations such as $\chi_T = \beta \left\langle \delta M^2 \right\rangle$ and $K_T = \beta \left\langle \delta V^2 \right\rangle / V$, in order to consider the existence of macrostates with anomalous values in response functions such as the isothermal susceptibility in a paramagnetic system $\chi_T = \partial M / \partial H$ or the isothermal compressibility of a fluid system $K_T = -V^{-1} \left(\partial V / \partial p \right)_T$. Finally, we discuss some simple application examples and perform a numerical testing of some consequences of the present generalization by studying the anomalous diamagnetic states of the 2D Ising model through Monte Carlo simulations.

PACS numbers: 05.20.Gg; 05.40.-a; 75.40.-s; 02.70.Tt

[1]Departamento de Física, Universidad de Pinar del Río, Martí 270, Esq. 27 de Noviembre, Pinar del Río, Cuba.
[2]Departamento de Física, Universidad Católica del Norte, Av. Angamos 0610, Antofagasta, Chile.

1. Introduction

According to the well-known fluctuation relation:

$$C = k_B \beta^2 \left\langle \delta U^2 \right\rangle \tag{1}$$

between the heat capacity C and the energy fluctuations $\left\langle \delta U^2 \right\rangle$, the heat capacity should be *nonnegative*. However, such a conclusion is only an illusion. Since the first theoretical demonstration about the existence of macrostates with negative heat capacities $C < 0$ by Lyndel-Bell in the astrophysical context [1], this anomaly has been observed in several systems [2, 3, 4, 5, 6, 7, 8]. The fluctuation relation (1) directly follows from the consideration of the Gibbs' canonical ensemble:

$$dp_c \left(U \left| \beta \right. \right) = \frac{1}{Z \left(\beta \right)} \exp \left(-\beta U \right) \Omega \left(U \right) dU, \tag{2}$$

which accounts for the equilibrium thermodynamic properties of a system in thermal contact with a heat bath at constant temperature T when other thermodynamical variables like the system volume V or a magnetic field H are kept fixed, where $\beta = 1/k_B T$. As elsewhere shown [8, 9], macrostates with $C < 0$ can exist within the microcanonical description of a given system, but these states are *thermodynamically* *unstable* under the external influence imposed to the system within the Gibbs' canonical ensemble.

A suitable extension of this standard result has been recently derived [10]:

$$C = k_B \beta^2 \left\langle \delta U^2 \right\rangle + C \left\langle \delta \beta^\omega \delta U \right\rangle , \tag{3}$$

which considers a *system-surroundings* equilibrium situation where the inverse temperature β^ω of a given thermostat exhibits correlated fluctuations with the total energy U of the system under study as a consequence of the underlying thermodynamic interaction. This last generalization differs from the canonical equilibrium situation in the fact that the internal thermodynamical state of the thermostat can be affected by the presence of the system under study, constituting in this way a more general framework than the usual canonical ($\delta\beta^\omega = 0$) and microcanonical ensembles ($\delta U = 0$).

The new fluctuation relation (3) defines a criterion capable of detecting the presence of a regime with $C < 0$ in the microcanonical caloric curve $1/T\left(U\right) = \partial S\left(U\right)/\partial U$ of the system through the correlated fluctuations of the inverse temperature β^ω and the energy U of the system itself. In fact, it asserts that macrostates with $C < 0$ are thermodynamically stable provided that the influence of the thermostat obeyed the inequality $\left\langle \delta\beta^\omega \delta U \right\rangle > 1$. Consequently, any attempt to impose the canonical conditions $\delta\beta^\omega \rightarrow 0$ leads to very large energy fluctuations $\delta U \rightarrow \infty$ that induce the thermodynamic instability of such anomalous macrostates.

As already discussed in a previous work [11], the fluctuation-dissipation relation (3) has interesting connections with different questions within statistical mechanics, such as the extension of canonical Monte Carlo methods to allow the study macrostates with negative heat capacities and to avoid the super-critical slowing down of first-order phase transitions [12], the justification of a complementary relation between energy and temperature [13, 14, 15, 16, 17, 18, 19, 20, 21], the development of a geometric formulation for fluctuation theory [22, 23], as well as the Mandelbrot's derivation of statistical mechanics from inference theory [16]. However, Eq.(3) is applicable to those equilibrium situations where there is only involved the conjugated pair energy-temperature. Thus, this result merely constitutes a first step towards the development of an extension of *equilibrium fluctuation-dissipation relations* compatible with the existence of anomalous response functions [27, 28, 29, 30, 31, 32].

In this paper, we shall extend our previous results to an equilibrium situation with several control parameters, that is, we shall develop a special approach of fluctuation theory [24] in order to arrive at a suitable generalization of the familiar fluctuation relations:

$$C_V = \beta^2 \left\langle \delta Q^2 \right\rangle , \; V K_T = \beta \left\langle \delta V^2 \right\rangle , \; \chi_T = \beta \left\langle \delta M^2 \right\rangle , \tag{4}$$

compatible with the existence of macrostates with *anomalous values in response functions*, such as negative heat capacities $C_V < 0$, isothermal compressibilities $K_T < 0$ or isothermal magnetic susceptibilities $\chi_T < 0$ in those magnetic systems where this response function is expected to be nonnegative.

2. The proposal

2.1. Notation conventions

From the standard perspective of statistical mechanics, a system-surroundings equilibrium situation with several control parameters is customarily described by using the *Boltzmann-Gibbs distributions* [24]:

$$dp_{BG}\left(U,X\right|\beta,Y)=\frac{1}{Z\left(\beta,Y\right)}\exp\left[-\beta\left(U+YX\right)\right]\Omega\left(U,X\right)dUdX. \tag{5}$$

The quantities $X=(V,M,P,N_i,\ldots)$ represent other macroscopic observables acting in a given application (generalized displacements) such as the volume V, the magnetization M and polarization P, the number of chemical species N_i, etc.; with $Y=(p,-H,-E,-\mu_i,\ldots)$ being the corresponding conjugated thermodynamic parameters (generalized forces): the external pressure p, magnetic H and electric E fields, the chemical potentials μ_i, etc. In the sake of simplicity, we shall consider the Boltzmann's constant $k_B\equiv 1$ along this work, so that, β in Eq.(5) represents the inverse temperature, $\beta=1/T$.

Conventionally, the system energy U is the most important physical observable in thermodynamics and statistical mechanics, and hence, it is always distinguished from the other macroscopic observables X. However, we shall not start from this distinction in the present approach, a consideration that allows us to deal with a more simple notation in our analysis. Let us consider the following convention for the physical observables $(U,X)\rightarrow I=(I^1,I^2,\ldots)$ and $(\beta,Y)\rightarrow\beta=(\beta_1,\beta_2,\ldots)$ for the thermodynamic parameters, in a way that the total differential of entropy S can be rewritten as follows: $dS=\beta\left(dU+YdX\right)\rightarrow dS=\beta_1 dI^1+\beta_2 dI^2+\ldots\equiv\beta_i dI^i$. We will use hereafter the Einstein's summation convention, which allows to rewrite the probabilistic weight of the Boltzmann-Gibbs distribution (5) as:

$$\omega_{BG}\left(I\right|\beta)=\frac{1}{Z\left(\beta\right)}\exp\left(-\beta_i I^i\right). \tag{6}$$

2.2. Starting considerations on the distribution function

At first glance, all that it is necessary to demand in order to extend fluctuation relations (4) is to start from a probabilistic distribution more general than the Boltzmann-Gibbs distributions (5). From a mathematical point of view, a distribution that seems to describe the thermodynamical behavior of a system under a general influence of an environment is the ansatz:

$$dp\left(I\right)=\omega\left(I\right)\Omega\left(I\right)dI, \tag{7}$$

whose generic probabilistic weight $\omega\left(I\right)$ admits the Boltzmann-Gibbs form (6) as a particular case. We shall shown in the next subsections that the hypothesis (7), although simple, is sufficiently general to achieve our purposes. In fact, this expression is just a direct extension of the ansatz used in our previous work [10]:

$$dp\left(U\right)=\omega\left(U\right)\Omega\left(U\right)dU \tag{8}$$

to a more general case with several macroscopic observables. Let us now discuss under what general background conditions the mathematical form (7) can be supported by physical arguments.

A particular and conventional situation is that where the system under study and its environment can be considered as two *separable and independent finite subsystems* that constitute a closed system in thermodynamic equilibrium. By admitting only additive physical observables (e.g.: energy, volume, particles number or electric charge) obeying the constraint $I_T = I + I_A$, a simple ansatz for the distribution function is given by:

$$dp\left(I|\,I_T\right) = \frac{1}{W\left(I_T\right)}\Omega_e\left(I_T - I\right)\Omega\left(I\right)dI, \tag{9}$$

where $\Omega\left(I\right)$ and $\Omega_e\left(I_e\right)$ are the densities of states of the system and the environment, respectively, and $W\left(I_T\right)$, the partition function that ensures the normalization condition:

$$\int_\Gamma dp\left(I|\,I_T\right) = 1. \tag{10}$$

Here, Γ represents all admissible values of the system observables I.

Although Eq.(9) provides a simple interpretation for Eq.(7), it clearly dismisses some important practical situations where the *additive constraint* $I_T = I + I_e$ cannot be ensured for all physical observables involved in the system-environment thermodynamic interaction. Significant examples of *unconstrained observables* are the magnetization $\mathbf{M}$ or the electric polarization $\mathbf{P}$ associated with magnetic and electric systems respectively. A formal way to overcome such a difficulty is carried out by representing the density of state of the environment Ω_e in terms of the observables of the system I under study and the set of control parameters a of the given physical situation, $\Omega_e = \Omega_e\left(I|\,a\right)$. Such an explicit dependence of the density of states of the environment on the internal state of the system follows from their mutual interaction, which leads to the existence of *correlative effects* between these systems. Thus, one can express the corresponding distribution function as follows:

$$dp\left(I|\,a\right) = \frac{1}{W\left(a\right)}\Omega_e\left(I|\,a\right)\Omega\left(I\right)dI. \tag{11}$$

The subset Γ_a of all admissible values of the physical observables I now depend on the control parameters a.

Rigorously speaking, both *separability* and *additivity* represent suitable idealizations of the real physical conditions associated with the large thermodynamic systems with a microscopic dynamics driven by short-range interactions. These conditions cannot be naively extended to systems with long-range interactions, which constitute a relevant framework for the existence of macrostates with anomalous response functions. As already mentioned in the introductory section, remarkable examples are the astrophysical systems, whose thermodynamic description is hallmarked by the existence of *negative heat capacities* $C < 0$. Although one cannot trivially divide a long-range interacting system into independent subsystems, an *effective separability* can be arisen as a direct consequence of the underlying microscopic dynamics in some practical situations.

For example, it is physically admissible to speak about a separability between a globular cluster and its nearby galaxy despite of the long-range character of the gravitational interaction. Since separability is, at least, an important condition to attribute a reasonable physical meaning to the description of a system as an individual entity, we shall demand this property as necessary requirement for the applicability of the ansatz (7).

The existence of long-range interactions also implies the incidence of *long-range correlations*. This last feature, by itself, modifies in a significant way the thermodynamic behavior and the macroscopic dynamics of a physical system. If one considers the existing analogy with conventional systems at the *critical points* (where correlation length ξ diverges, $\xi \to \infty$), the dynamics of long-range interacting systems should undergo the existence of some collective phenomena with a very slow relaxation towards the final equilibrium [9], as example, the so-called *violent relaxation* that appears during the *collisionless dynamics* of the astrophysical systems [33]. Under this last circumstance, it is usual that a long-range interacting system cannot be found in a final thermodynamic equilibrium, but in a long-living *quasi-stationary state* [25, 26]. The above dynamical features establish certain analogies among the long-range interacting systems with other physical systems with a complex microscopic dynamics such as *turbulent fluids* [34] and *glassy systems* [35].

According to the above reasonings, one cannot expect that the probabilistic weight $\omega(I)$ in Eq.(7) can be always interpreted in terms of the density of states of the environment Ω_e as the cases associated with Eqs.(9) or (11), since they demand the existence of a final thermodynamic equilibrium and the non-incidence of long-range interactions. Certainly, few systems in Nature are in absolute and final equilibrium, since it presupposes that all radiative materials would decayed completely and the nuclear reactions would have transmuted all nuclei to the most stable of isotopes. Thus, the hypothesis about the existence of a metastable equilibrium could be sufficient for the applicability of thermodynamics in many practical situations. Hereafter, we shall assume that *the probabilistic weight $\omega(I)$ accounts for a general thermodynamic influence of an environment*, which is found, at least, in conditions of *meta-stability*. This type of *ad hoc* hypothesis, that is, the using of *generalized statistical ensemble* with the form (7), is usual in the statistical mechanics literature in the last decades. This kind of methodology has proved to be an useful alternative to describe thermodynamic features of systems with a complex microscopic dynamics. The reader can find an unifying framework of all these generalized distribution functions in the so-called *Superstatistics*, a theory recently proposed by C. Beck and E.G.D Cohen [36].

The specific mathematical form of the probabilistic weight $\omega(I)$ depends, in general, on the internal structure of the environment, the character of its own equilibrium conditions, and the nature of forces driving the underlying system-environment thermodynamic interaction. Fortunately, the exact mathematical form of the function $\omega(I)$ is unimportant in our subsequent development. Despite of its generality, we shall show that the internal thermodynamic state of the environment is fully characterized by a finite set of *control thermodynamic parameters* $\beta^{\omega} = (\beta^{\omega}_1, \beta^{\omega}_2, \ldots)$ that constitute a natural extension of the constant parameters $\beta=(\beta_1,\beta_2,\ldots)$ of the Boltzmann-Gibbs distributions (6). Such a reduced description is analogous to the equilibrium situation described within the Gibbs' canonical ensemble (2), where the external influence is determined by the thermostat temperature T regardless its internal structure and composition, as long as the thermostat exhibited a very large heat capacity and guaranteed a thermal contact with the system under analysis.

2.3. Thermodynamic control parameters of the environment

The direct way to arrive at a suitable extension of the thermodynamic control parameters $\beta=(\beta_1,\beta_2,\ldots)$ of the Boltzmann-Gibbs distributions (6) is achieved by appealing to the *conditions of thermodynamic equilibrium* such as the condition of *thermal equilibrium*, $T^{\omega}=T^{s}$, the condition of *mechanical equilibrium*, $p^{\omega}=p^{s}$, the condition of *chemical equilibrium*, $\mu^{\omega}=\mu^{s}$, etc. As usual, the conditions of thermodynamic equilibrium can be derived as the *stationary conditions* associated with the *most likely macrostate* $\bar{I}$ of the distribution function $\rho(I)=\omega(I)\Omega(I)$:

$$\sup_{\bar{I}}\left\{\omega(I)\Omega(I)\right\}\Rightarrow\frac{\partial}{\partial I^{i}}\log\left\{\omega\left(\bar{I}\right)\Omega\left(\bar{I}\right)\right\}=0, \tag{12}$$

which can be conveniently rewritten as follows:

$$\beta_i^{\omega}\left(\bar{I}\right)=\beta_i^{s}\left(\bar{I}\right). \tag{13}$$

Here, β_i^{s} are the system (microcanonical) thermodynamic parameters:

$$\beta_i^{s}(I)=\frac{\partial S(I)}{\partial I^{i}} \tag{14}$$

derived from the *coarsed grained* entropy $S(I)\equiv\log W(I)$ with $W(I)=\Omega(I)\,\delta c$ (δc is a small constant which makes the volume W dimensionless), while *the quantities* β_i^{ω} *defined by*:

$$\beta_i^{\omega}(I)=-\frac{\partial\log\omega(I)}{\partial I^{i}} \tag{15}$$

are the control thermodynamic parameters of the environment. This last definition constitutes a fundamental consideration in the present development.

As expected, the environment control thermodynamic parameters (15) drop to the constant parameters $\{\beta_i\}$ of the Boltzmann-Gibbs distributions (6) by replacing the general probabilistic weight $\omega(I)$ by the one associated with this last ensemble $\omega_{BG}(I)$. Similarly, they drop to its microcanonical definitions (14) when the environment is just a short-range interacting system in a final thermodynamic equilibrium, that is, when the probabilistic weight ω is proportional to the states density Ω_e. However, these thermodynamic parameters turn *effective* in the other cases, overall, in those practical situations where the probabilistic weight $\omega(I)$ corresponds to an environment in a *metastable or quasi-stationary equilibrium.* The possibility to extend this type of thermodynamic concepts to situations of metastability is not strange, since there exist robust evidences supporting the introduction of concept of *effective temperature* in non-equilibrium systems (see in ref.[35] and references therein).

A remarkable feature of the environment thermodynamic parameters (15) is *their implicit dependence on the instantaneous macrostate I of the system*, since these thermodynamic quantities are only constant for the particular case of the Boltzmann-Gibbs distributions. Consequently, it is almost a rule rather than an exception that *the underlying system-environment interaction provokes the existence of non-vanishing correlated fluctuations among the environment thermodynamic parameters and system observables*, $\langle \delta\beta_i^{\omega}\delta I^j\rangle \neq 0$. Such a realistic behavior of the interacting macroscopic systems is systematically disregarded by the use of Boltzmann-Gibbs distributions (6).

The quantities $\{\beta_i^{\omega}\}$ act as control parameters during the system-environment thermodynamic interaction. A simple way to illustrate this role is by using the probabilistic weight $\omega\left(I\right)$ in the Metropolis Monte Carlo simulation [37] of the system under the present equilibrium conditions. The acceptance probability for a Metropolis move $p\left(I\right|I+\Delta I)$ is given by:

$$p\left(I\right|I+\Delta I)=\min\left\{\frac{\omega\left(I+\Delta I\right)}{\omega\left(I\right)},1\right\}. \tag{16}$$

The fluctuations of the macroscopic observables satisfy the condition $|\Delta I|\ll|I|$ whenever the size N of the system is sufficiently large, allowing in this way to introduce the following approximation:

$$\log\left\{\frac{\omega\left(I+\Delta I\right)}{\omega\left(I\right)}\right\}\simeq\frac{\partial\log\omega\left(I\right)}{\partial I^i}\cdot\Delta I^i \tag{17}$$

and rewrite the acceptance probability (16) in an analogous fashion to the usual Metropolis algorithm based on the Boltzmann-Gibbs weight (6):

$$p\left(I\right|I+\Delta I)\simeq\left\{\exp\left[-\beta_i^{\omega}\left(I\right)\Delta I^i\right],1\right\}. \tag{18}$$

This last expression shows that the environment thermodynamic parameters (15) provide a simple extension of control parameters of Boltzmann-Gibbs distribution (6). Latter, we shall employed this acceptance probability to perform numerical simulations.

2.4. Derivation of some general fluctuation relations

As already commented, the set Γ represents all admissible values of the system observables $I=(I^2,I^2,\dots I^n)$. Let us denote by $\partial_i A\left(I\right)=\partial A\left(I\right)/\partial I^i$ the first partial derivatives of a given real function $A\left(I\right)$ defined on Γ. In general, the distribution function $\rho(I)=\omega(I)\Omega(I)$ of Eq.(7) should obey some mathematical properties associated with its physical significance. The are the following:

C1. *Existence*: The distribution function ρ is a nonnegative, bounded, continuous and differentiable function on Γ.

C2. *Normalization*: The distribution function ρ obeys the normalization condition:

$$\int_{\Gamma}\rho\left(I\right)dI=1. \tag{19}$$

C3. *Boundary conditions*: The distribution function ρ vanishes with its first partial derivatives $\{\partial_i \rho\}$ on the boundary $\partial\Gamma$ of the set Γ. Moreover, the distribution function ρ satisfies for $\alpha \leq n+1$ the condition:

$$\lim_{|I|\to\infty} |I|^{\alpha} \rho(I) = 0, \tag{20}$$

when the boundary $\partial\Gamma$ contains the infinite point $\{\infty\}$ of R^n.

C4. *Existence of the second partial derivatives*: The distribution function ρ admits second partial derivatives $\{\partial_i \partial_j \rho\}$ almost everywhere on Γ, with the exception of a certain subset $\Sigma \subset \Gamma$ with vanishing measure $\mu(\Sigma)$:

$$\mu(\Sigma) = \int_{\Sigma} \rho(I)\, dI = 0. \tag{21}$$

The mathematical conditions (C1-C3) constitute a direct extension of conditions (C1-C3) of our previous work to a case with several system observables (see in subsection 3.3 of Ref.[10]). Now, we have modified condition (C3) with the inclusion of the property (20), as well as we have considered a fourth requirement, the condition (C4). In general, conditions (C3) and (C4) ensure the existence of the expectation values $\langle I^i \rangle$ and some relevant correlations functions such as $\langle \delta I^i \delta I^j \rangle$.

Let us now introduce the quantity $\eta_i = \eta_i(I) = \beta_i^{\omega}(I) - \beta_i^{s}(I)$ defined as the difference between the i-th thermodynamic parameters of the environment and the system, respectively. By taking into consideration definitions (14) and (15), the thermodynamic quantity η_i can be associated with a *differential operator* $\hat{\eta}_i$ defined by:

$$\hat{\eta}_i = -\partial_i \tag{22}$$

due to the validity of the mathematical identity:

$$\hat{\eta}_i \rho(I) \equiv \eta_i \rho(I). \tag{23}$$

Its demonstration reads as follows: the partial first derivative of ρ can be rewritten as $-\partial_i \rho = (-\partial_i \log \rho)\rho$ and $-\partial_i \log \rho \equiv -\partial_i \log \omega - \partial_i S = \beta_i^{\omega} - \beta_i^{s} = \eta_i$, with $S = \log W$ being the coarsed grained entropy with $W = \Omega \delta c$.

The set of operators $\{\hat{\eta}_i\}$ satisfy the following identities:

$$[A(I), \hat{\eta}_i] = \partial_i A(I), \quad [\hat{\eta}_i \hat{\eta}_j] = 0, \tag{24}$$

with $\left[\hat{A}, \hat{B}\right] = \hat{A}\hat{B} - \hat{B}\hat{A}$ being the so-called *commutator* between two operators $\hat{A}$ and $\hat{B}$. A remarkable particular case of the first commutator is obtained when $A = \hat{I}^j \equiv I^j$:

$$\left[\hat{I}^j, \hat{\eta}_i\right] = \delta_i^j. \tag{25}$$

The consecutive application of two operators $\hat{\eta}_i$ on the distribution function ρ leads to the following result:

$$\hat{\eta}_i \hat{\eta}_j \rho = \hat{\eta}_i (\hat{\eta}_j \rho) = \hat{\eta}_i (\eta_j \rho) = (-\partial_i \eta_j + \eta_i \eta_j)\rho. \tag{26}$$

Notice here that condition (C4) is crucial for the existence of the partial derivatives $\partial_i \eta_j$.

Now, let us combine the general mathematical properties (C1-C4) of the distribution function $\rho(I)$ with the operational representation (23) of the thermodynamic quantity η_i to demonstrate the validity of the following expectation values:

$$\langle \eta_i \rangle = 0, \ \ \left\langle \eta_i I^j \right\rangle = \delta_i^j, \ \ \langle -\partial_i \eta_j + \eta_i \eta_j \rangle = 0, \tag{27}$$

which arise as particular cases of the following identity:

$$\langle \partial_i A \rangle = \langle A \eta_i \rangle \tag{28}$$

when $A = 1$, $A = I^j$ and $A = \eta_j$, respectively. Here, $A(I)$ is a differentiable function that satisfies the condition $A(I) \leq \lambda \left|I\right|^2$ when $|I| \to \infty$ if the set Γ is not finite.

The demonstration start from the consideration of a set of arbitrary constants $\{a^i\}$, which is employed to rephrase the integral:

$$a^i \langle A\eta_i \rangle = \int_\Gamma A a^i \eta_i \rho dI \tag{29}$$

by using a vector $\vec{J}$ with components $J^i = a^i \rho$ as follows:

$$\int_\Gamma A a^i \eta_i \rho dI = \int_\Gamma A a^i \hat{\eta}_i \rho dI = -\int_\Gamma A \vec{\nabla} \cdot \vec{J} dI. \tag{30}$$

By using the identity:

$$A \vec{\nabla} \cdot \vec{J} = -\vec{\nabla} \cdot \left(A\vec{J} \right) + \vec{J} \cdot \vec{\nabla} A, \tag{31}$$

the integral (30) can be rewritten as follows:

$$-\int_\Gamma A \vec{\nabla} \cdot \vec{J} dI = -\int_{\partial\Gamma} A\vec{J} \cdot d\vec{\sigma} + \int_\Gamma \vec{J} \cdot \vec{\nabla} A dI. \tag{32}$$

The surface integral over the boundary $\partial\Gamma$ vanishes as a consequence of condition (C3). This conclusion is evident when the set Γ is finite since $J|_{\partial\Gamma} = 0$. Conversely, this vanishing takes place as a consequence of the limit:

$$\lim_{|I| \to \infty} |I|^{n-1} \left| A(I) \rho(I) \right| \leq \lim_{|I| \to \infty} \lambda \left|I\right|^{n+1} \left|\rho(I)\right| = 0. \tag{33}$$

Identity (28) is obtained by considering $a^j = 0$ for $j \neq i$ and $a^i = 1$ in the final expression:

$$a^j \langle A \eta_j \rangle = \int_\Gamma \vec{J} \cdot \vec{\nabla} A dI \equiv a^j \langle \partial_j A \rangle. \tag{34}$$

By taking into account the vanishing of $\langle \eta_i \rangle$ and well as the definition of the correlation function $\langle \delta\eta_i \delta I^j \rangle \equiv \langle \eta_i I^j \rangle - \langle \eta_i \rangle \langle I^j \rangle$, the second relation of Eq.(27) can be conveniently rewritten as follows:

$$\left\langle \delta\eta_i \delta I^j \right\rangle = \delta_i^j, \tag{35}$$

which is hereafter referred to as the *fundamental fluctuation relation.* By using this same procedure, the third relation of Eq.(27) can be rephrased as follows:

$$\langle \partial_i \eta_j \rangle = \langle \delta\eta_i \delta\eta_j \rangle, \tag{36}$$

which is hereafter referred to as the *associated fluctuation relation.* Let us now analyze the physical consequences of these mathematical results.

The vanishing of expectation values $\langle\eta_i\rangle$ in Eq.(27) drops to the following thermodynamic relations:

$$\langle\beta_i^{\omega}\rangle = \langle\beta_i^{s}\rangle\,, \tag{37}$$

whose mathematical form explains by itself that the *conditions of thermodynamic equilibrium* are not only supported by the argument of *the most likely macrostate* (see Eq.(13) in the previous subsection), but it can be rigorous expressed in terms of statistical expectation values. In fact, the present version (37) of the thermodynamic equilibrium conditions has a more general applicability than the conventional version (13), since the interpretation of the state of equilibrium through the most likely macrostate $\bar{I}$ is physically relevant for the large thermodynamic systems where thermal fluctuations can be disregarded.

Conventionally, pairs of thermodynamic quantities such as energy U and temperature T, volume V and pressure p, magnetization $\mathbf{M}$ and magnetic field $\mathbf{H}$, etc., are regarded as *conjugated thermodynamic quantities.* In this framework, commutative relation (25) allows to refer to the pairs (I^i,η_i) as *thermodynamic complementary quantities.* Examples of thermodynamic complementary quantities are the energy U and the inverse temperature difference $\eta_U = 1/T^{\omega} - 1/T^{s}$ between the environment (ω) and system (s), or the generalized displacement X and the quantity $\eta_X = Y^{\omega}/T^{\omega} - Y^{s}/T^{s}$ defined in terms of the generalized forces (Y^s, Y^{ω}) of the system and the environment respectively. The physical relevance of this new conceptualization relies on the fact that *the thermal fluctuations of thermodynamic complementary quantities cannot be simultaneously reduced to zero.* In order to shown this claim, let us first notice by considering Eq.(35) that only the fluctuations corresponding to thermodynamic complementary quantities (I^i,η_i) are *statistically dependent.* By rewriting the fluctuation relations of complementary quantities through the known *Schwarz inequality*:

$$\left\langle\delta A^2\right\rangle\left\langle\delta B^2\right\rangle \geq \left\langle\delta A\delta B\right\rangle^2, \tag{38}$$

one obtains the following remarkable relations:

$$\Delta\eta_i\Delta I^i \geq 1, \tag{39}$$

where $\Delta x \equiv \sqrt{\langle\delta x^2\rangle}$ denotes the thermal dispersion of the quantity x. These inequalities obey the form of *uncertainty relations* rather analogous to the ones associated with quantum theories. In fact, Eq.(39) is a direct extension of the energy-temperature uncertainty relation obtained in our previous work [10]:

$$\Delta U\Delta(1/T^{\omega} - 1/T^{s}) \geq 1 \tag{40}$$

to other conjugated thermodynamic quantities such as the generalized displacement X and the generalized force Y:

$$\Delta X\Delta(Y^{\omega}/T^{\omega} - Y^{s}/T^{s}) \geq 1. \tag{41}$$

By generalizing our previous result, the physical significance of thermodynamic uncertainty relation (39) is the following:

Claim 1: *It is impossible to perform a simultaneous exact determination of the system conjugated thermodynamic parameter I^{i} and $\beta_{i}^{s}=\partial_{i}S$ by using an experimental procedure based on the thermal equilibrium condition (37) with a second system regarded here as a "measuring apparatus": any attempt to reduce to zero the thermal uncertainties of the differences between their thermodynamic parameters $\Delta\eta_{i}\rightarrow 0$ involves a strong perturbation on the conjugated macroscopic observable $\Delta I^{i}\rightarrow\infty$, and hence, I^{i} becomes indeterminable; any attempt to reduce this perturbation to zero, $\Delta I^{i}\rightarrow 0$, makes impossible to determine the conjugated thermodynamic parameter β_{i}^{s} by using the condition of thermal equilibrium (37) since $\Delta\eta_{i}\rightarrow\infty$.*

The present result constitutes a clear evidence about the existence of *complementary relations in any physical theory with a statistical formulation*, an idea postulated by Bohr in the early days of quantum mechanics [13, 14] with a long history in literature [15, 16, 17, 18, 19, 20, 21]. Although our results are intimately related with other approaches based on fluctuation theory [15, 17, 20], there exist subtle but fundamental differences distinguishing the present proposal from other attempts carried out in the past. The interested reader can see a more complete discussion about this subject in Ref.[38].

Let us now discuss some consequences of the associated fluctuation relation (36). The quantities ζ_{ij}:

$$\zeta_{ij}=\langle\partial_{i}\eta_{j}\rangle=-\int_{\Gamma}\rho\left(I\right)\partial_{i}\partial_{j}\log\rho\left(I\right)dI \tag{42}$$

are just the expectation values of the Hessian of the logarithm of the distribution function $P_{ij}\left(I\right)$:

$$P_{ij}\left(I\right)=-\partial_{i}\partial_{j}\log\rho\left(I\right), \tag{43}$$

which can be referred to as the *probabilistic Hessian.* Due to the positive definite character of the correlation matrix $O=\left[O_{ij}\right]$ with components $O_{ij}=\langle\delta\eta_{i}\delta\eta_{j}\rangle$:

$$a^{i}a^{j}O_{ij}\equiv\left\langle\left(a^{i}\delta\eta_{i}\right)^{2}\right\rangle\geq 0, \tag{44}$$

as well as the applicability of the associated fluctuation relation (36), $\zeta_{ij}=O_{ij}$, the quantities ζ_{ij} constitute the components of *a positive definite symmetric matrix* $\zeta=\left[\zeta_{ij}\right]$.

The positive definite character of matrix $\left[\zeta_{ij}\right]$ does not imply a positive definite character of the probabilistic Hessian (43) for any macrostate $I\in\Gamma$. By admitting the existence of a certain region $\mathcal{A}^{\omega}\subset\Gamma$ where this last condition does not hold for any admissible value of the internal control parameters that drive the form of the probabilistic weight $\omega(I)$‡, the positive definite character of ζ_{ij} indicates that *macrostates belonging to the subset $\mathcal{A}^{\omega}$ have always a poor contribution to the integral of* Eq.(42). If the system size N is sufficiently large, the thermal fluctuations δI^{i} and

---

‡ For example, the thermodynamic parameters (β, Y) of the Boltzmann-Gibbs distribution (5), the total additive quantities I_{T} of distribution (9), or the formal parameters a of distribution (11.)

$\delta\eta_i$ turn negligible and the distribution function $\rho(I)$ significantly differs from zero in a very small neighborhood around its local maxima. In this limit, all system macrostates $I \in \mathcal{A}^{\omega}$ turns *inaccessible or thermodynamically unstable.* Since the probabilistic Hessian (43) depends on the particular mathematical form of the probabilistic weight $\omega(I)$ that accounts for the external influence of the environment, it is possible to claim the following:

Claim 2: *The thermodynamic stability conditions and the fluctuating behavior of a given open system in (a quasi-stationary) equilibrium depend on the nature of external conditions which have been imposed to.*

This last conclusion appears to be self-evident and natural, but it contracts with some familiar equilibrium fluctuation relations such as the ones shown in Eq.(4), which seem to suggest (falsely) that system stability and fluctuating behavior only depend on its intrinsic properties: *the response functions.* However, such a conclusion is strictly correct for an equilibrium situation described by the Boltzmann-Gibbs distributions (6). An illustrative example is the framework of application of the canonical fluctuation relation $C = \beta^2 \langle \delta U^2 \rangle$. The internal energy U fluctuates when the system is put in a thermal contact with a bath, but it remains constant when the system is put in an energetic isolation, and hence, there is no connection between the energy fluctuations and the heat capacity for the second situation.

2.5. Gaussian approximation

Fluctuation relations (35) and (36) constitute two general mathematical results. The price to pay in order to arrive at a suitable extension of the familiar equilibrium fluctuation relations (4) is the consideration of a *Gaussian approximation* for thermal fluctuations, which appears as a licit treatment when the size N of the system under consideration is sufficiently large.

Let us admit that the distribution function (7) only exhibits one (Gaussian) peak around the most likely macrostate $\bar{I}$:

$$\rho(I) \propto \exp\left[-\frac{1}{2}C_{ij}\delta I^i \delta I^j\right], \tag{45}$$

with $\delta I^i = I^i - \bar{I}^i$. Since the thermal fluctuations δI are small, the fluctuations of any continuous and differentiable function $A(I)$ can be expressed in a first-order approximation as follows:

$$\delta A(I) \simeq \partial_i A\left(\bar{I}\right)\delta I^i. \tag{46}$$

Formally, the thermodynamic parameters of the system $\beta^s_i = \partial_i S(I)$ depend on the macroscopic observables I. Thus, their fluctuations $\delta\beta^s_i$ can be expressed by using the fist-order approximation (46):

$$\delta\beta^s_i = \partial_k \beta^s_i\left(\bar{I}\right)\delta I^k = H_{ik}\left(\bar{I}\right)\delta I^k, \tag{47}$$

with $H_{ij}\left(\bar{I}\right)=\partial_i\partial_j S\left(\bar{I}\right)$ being the value of the *entropy Hessian* at the most likely macrostate. The substitution of Eq.(47) into the fundamental fluctuation relation (35) allows to obtain the following expression:

$$\left\langle\delta\beta_i^{\omega}\delta I^j\right\rangle=\delta_i^j+H_{ik}\left(\bar{I}\right)\left\langle\delta I^k\delta I^j\right\rangle. \tag{48}$$

By using these same reasonings, the associated fluctuation relation (36) can be rewritten as follows:

$$\partial_i\eta_j\left(\bar{I}\right)=\left\langle\delta\eta_i\delta\eta_j\right\rangle. \tag{49}$$

Result (49) evidences that the regions of thermodynamic stability are those macrostates where the probabilistic Hessian:

$$P_{ij}\left(\bar{I}\right)=\partial_i\eta_j\left(\bar{I}\right)=-\partial_i\partial_j\left\{\log\omega\left(\bar{I}\right)+S\left(\bar{I}\right)\right\}, \tag{50}$$

is a positive definite matrix. Besides, Eqs.(48) and (49) indicate that the system fluctuating behavior and its stability conditions depend on the external conditions which have been imposed to (Claim 2), which are characterized here by the correlations $\langle\delta\beta_i^{\omega}\delta I^j\rangle$ and the Hessian of the logarithm of probabilistic weight with opposite sign $F_{ij}\left(\bar{I}\right)=-\partial_i\partial_j\log\omega\left(\bar{I}\right)$.

It is not difficult to verify that these identities provide the same qualitative information about the system thermodynamic stability. For example, the equilibrium conditions associated with the Boltzmann - Gibbs distributions (6) follows from the imposition of constraint $\delta\beta_i^{\omega}=0$ into Eq.(48). Thus, the correlation matrix $C^{ij}=\left\langle\delta I^k\delta I^j\right\rangle$ is given by the inverse matrix of the entropy Hessian $H_{ij}\left(\bar{I}\right)$ with opposite sign:

$$-H^{ij}\left(\bar{I}\right)\equiv C^{ij}, \tag{51}$$

which is a well-known result of classical fluctuation theory [24]. Since the correlation function C^{ij} is always a positive definite matrix:

$$a_i a_j C^{ij}\equiv\left\langle\left(a_i\delta I^i\right)^2\right\rangle\geq 0, \tag{52}$$

the peaks of the Boltzmann-Gibbs distribution (5) will never be located in those regions where $H_{ij}\left(I\right)$ is a nonnegative definite matrix, that is, where the entropy S *is not a locally concave function*. Thus, these macrostates turn inaccessible in the limit of large system size N, so that, all them constitute the unstable region $\mathcal{A}^{\omega}=\mathcal{A}^{BG}$ of this particular ensemble. Clearly, this is the same conclusion derived from Eqs.(49) and (50). As a by-product of the present reasonings, we have arrive at the following:

Claim 3: *The direct observation of unstable macrostates* $I\in\mathcal{A}^{BG}$ *of the Boltzmann-Gibbs distributions (5) is only possible when* $\langle\delta\beta_i^{\omega}\delta I^j\rangle\neq 0$*, that is, by considering an equilibrium situation where the internal thermodynamic state of the environment is affected by thermodynamic interaction with the system.*

There exist within the Gaussian approximation a fluctuation relation which has no counterpart within the Boltzmann-Gibbs distributions since it involves the correlation matrix $B_{ij}=\left\langle\delta\beta_i^{\omega}\delta\beta_j^{\omega}\right\rangle$. By using the first-order approximation:

$$\delta\beta_i^{\omega}=\partial_j\beta_i^{\omega}\left(\bar{I}\right)\delta I^j=F_{ij}\left(\bar{I}\right)\delta I^j \tag{53}$$

one can express the correlation matrix $B_{ij}=\left\langle \delta\beta^{\omega}_{i}\delta\beta^{\omega}_{j}\right\rangle$ as follows:

$$\left\langle \delta\beta^{\omega}_{i}\delta\beta^{\omega}_{j}\right\rangle = F_{jk}\left(\bar{I}\right)\left\langle \delta\beta^{\omega}_{i}\delta I^{k}\right\rangle . \tag{54}$$

In the sake of simplicity, let us adopt the notation $\bar{F}_{ij}=F_{ij}\left(\bar{I}\right)$ and $\bar{H}_{ij}=H_{ij}\left(\bar{I}\right)$. By applying two times the fluctuation relation (48) and the approximation

$$\left\langle \delta\beta^{\omega}_{i}\delta I^{j}\right\rangle = \bar{F}_{im}\left\langle \delta I^{m}\delta I^{j}\right\rangle , \tag{55}$$

one obtains:

$$\left\langle \delta\beta^{\omega}_{i}\delta\beta^{\omega}_{j}\right\rangle = \bar{F}_{ij}+\bar{H}_{ij}+\bar{H}_{im}C^{mn}\bar{H}_{nj}. \tag{56}$$

This result can be conveniently rewritten by performing the matrix product with the correlation matrix $C^{ij}=\langle \delta I^{i}\delta I^{j}\rangle$:

$$\begin{aligned} B_{ik}C^{kj} &= \bar{F}_{ik}C^{kj}+\bar{H}_{ik}C^{kj}+\bar{H}_{im}C^{mn}\bar{H}_{nk}C^{kj} \\ &= \delta^{j}_{i}+2\bar{H}_{ik}C^{kj}+\bar{H}_{im}C^{mn}\bar{H}_{nk}C^{kj}. \end{aligned} \tag{57}$$

It is possible to recognize that the right hand side of the above equation is simply the term $M^{k}_{i}M^{j}_{k}$, where $M^{j}_{i}=\delta^{j}_{i}+\bar{H}_{ik}C^{kj}$ denotes the correlation matrix $M^{j}_{i}=\langle \delta\beta^{\omega}_{i}\delta I^{j}\rangle$ of Eq.(48). Thus, we have derived a *complementary fluctuation relation*:

$$\langle \delta\beta^{\omega}_{i}\delta\beta^{\omega}_{k}\rangle\left\langle \delta I^{k}\delta I^{j}\right\rangle = \left\langle \delta\beta^{\omega}_{i}\delta I^{k}\right\rangle\left\langle \delta\beta^{\omega}_{k}\delta I^{j}\right\rangle , \tag{58}$$

or its equivalent representation in matrix form:

$$B_{ik}C^{kj}=M^{k}_{i}M^{j}_{k}. \tag{59}$$

General speaking, the elements of the correlation matrix $M^{j}_{i}=\langle \delta\beta^{\omega}_{i}\delta I^{j}\rangle$ could be very difficult to measure in a given experimental situation, since it demands to perform simultaneous measurements of the system observables I and the thermodynamic parameters of the environment β^{e}. The complementary fluctuation relation (59) allows to obtain in an indirect way these fluctuating quantities by performing independent measurements of the fluctuations of the system observables I and the environment thermodynamic parameters β^{e}.

2.6. Extended equilibrium fluctuation-dissipation theorem

For convenience, let us re-derive here the fluctuation relations associated with the Boltzmann-Gibbs distribution (6). The direct way to carry out this aim is by starting from the partition function $Z\left(\beta\right)$:

$$Z\left(\beta\right)=\int_{\Gamma}\exp\left(-\beta_{i}I^{i}\right)\Omega\left(I\right)dI. \tag{60}$$

Let us denote by $\partial^{i}A\left(\beta\right)=\partial A\left(\beta\right)/\partial\beta_{i}$ the first partial derivatives of an arbitrary function $A\left(\beta\right)$ defined on the thermodynamic parameters β. It is easy to verify the validity of the following identities:

$$\left\langle I^{i}\right\rangle = -\partial^{i}\log Z\left(\beta\right),\ \ \left\langle \delta I^{i}\delta I^{j}\right\rangle = \partial^{i}\partial^{j}\log Z\left(\beta\right). \tag{61}$$

The substitution of the first relation into the second one yields:

$$\chi_{BG}^{ij} = C^{ij} = \left\langle \delta I^{i} \delta I^{j} \right\rangle , \tag{62}$$

where the symmetric matrix χ_{BG}^{ij}:

$$\chi_{BG}^{ij} = -\frac{\partial \left\langle I^{j} \right\rangle}{\partial \beta_{i}} \tag{63}$$

will be referred to as the *canonical response matrix*. Since the correlation matrix $C^{ij} = \langle \delta I^{i} \delta I^{j} \rangle$ is a positive definite matrix, the canonical response matrix χ_{BG}^{ij} is also a positive definite matrix.

The identity (62) is a rigorous result within the Boltzmann-Gibbs distributions (6). Notice that an analogous expression can be derived within the Gaussian approximation from the fluctuation relation (48) after imposing the restriction $\delta\beta_{i}^{\omega} = 0$, that is, Eq.(51), which can be rephrased as:

$$\chi_{s}^{ij} = \left\langle \delta I^{i} \delta I^{j} \right\rangle , \tag{64}$$

with χ_{s}^{ij} being the inverse matrix of the entropy Hessian with opposite sign evaluated at the most likely macrostate:

$$\chi_{s}^{ij} = -H^{ij} \left(\bar{I} \right) . \tag{65}$$

Comparison between Eqs.(62) and (64) leads to establish the relationship $\chi_{BG}^{ij} \sim \chi_{s}^{ij}$, which allows to refer to χ_{s}^{ij} as the *microcanonical response matrix* of the system.

The microcanonical response matrix χ_{s}^{ij} defined above is a *purely microcanonical quantity*, that is, it is an intrinsic thermodynamic quantity that can be attributed to an isolated system. Actually, the physical meaning of such a response matrix is *unclear*. In general, a response function characterizes the *sensibility of the system under the external control*, and consequently, such a concept is only relevant for an open system. While such a physical relevance is satisfied by the canonical response matrix χ_{BG}^{ij}, its applicability is only restricted to the case when the thermodynamic influence of the environment corresponds to the Boltzmann-Gibbs distributions (6), where the thermodynamic parameters $\{\beta_{i}\}$ are constant parameters instead of *fluctuating quantities*.

According to these last reasonings, the concept of response matrix χ^{ij} should be extended in order to be also applicable in a more general system-environment equilibrium situation. Since the physical observables I and the (control) thermodynamic variables β^{ω} undergoing, in general, thermal fluctuations, a simple operational definition for the concept of response matrix χ_{ω}^{ij} could be introduced in terms of expectation values as follows:

$$\chi_{\omega}^{ij} = -\frac{\partial \left\langle I^{j} \right\rangle}{\partial \left\langle \beta_{i}^{\omega} \right\rangle} \equiv -\frac{\partial \left\langle I^{j} \right\rangle}{\partial \left\langle \beta_{i}^{s} \right\rangle} . \tag{66}$$

Here, we have took into account the equilibrium conditions $\langle \beta_{i}^{\omega} \rangle = \langle \beta_{i}^{s} \rangle$.

Clearly, definition (66) contains the canonical response matrix χ_{BG}^{ij} as a particular case when $\delta\beta_{i}^{\omega} = 0 \Rightarrow \beta_{i}^{\omega} = \beta_{i} \sim$ *const*. The response matrix χ_{ω}^{ij} defined in this way is more realistic and general than the one associated with the canonical response matrix

(63). The canonical response matrix χ^{ij}_{BG} presupposes that the control parameters $\{\beta_i\}$ have no fluctuations. Such an exigency is clearly an idealization since *any real measuring process always involves a perturbation of the thermodynamic state of the measuring apparatus.* For example, both the temperature indicated by a thermometer and the pressure indicated by a barometer are affected by the thermodynamic influence of the system under study. In practice, the true values of temperature T and pressure p employed to calculate the heat capacity C_p at constant pressure and the isothermal compressibility K_T of a given system are actually *expectation values*, $\langle T\rangle$ and $\langle p\rangle$. Therefore, the practical operational definition of these concepts reads as follows:

$$C_p = \langle T\rangle \left(\frac{\partial \langle S\rangle}{\partial \langle T\rangle}\right)_{\langle p\rangle}, \; K_T = -\frac{1}{\langle V\rangle}\left(\frac{\partial \langle V\rangle}{\partial \langle p\rangle}\right)_{\langle T\rangle}, \tag{67}$$

where we have taken into consideration the stochastic character of the energy U, the volume V and the entropy $S = S(U, V)$.

As already discussed, the expectation values $\langle I^i\rangle$ and $\langle \beta^s_i\rangle$ are given within the Gaussian approximation by the corresponding values at the most likely macrostate $\bar{I}$, $\langle I^i\rangle = \bar{I}^i$ and $\langle \beta^s_i\rangle = \bar{\beta}^s_i = \beta^s_i\left(\bar{I}\right)$, which allows to rewrite Eq.(66) as follows:

$$\chi^{ij}_\omega = -\frac{\partial \bar{I}^j}{\partial \bar{\beta}^s_i} = \chi^{ij}_s\left(\bar{I}\right). \tag{68}$$

Claim 4: *The response matrix* χ^{ij}_ω *of Eq.*(66) *could be identified with the microcanonical response matrix* χ^{ij}_s (65) *at the most likely macrostate* $\bar{I}$ *as long as the Gaussian approximation applied.*

Equilibrium fluctuation relation (64) is inapplicable within the anomalous region $\mathcal{A}^{BG}$ of the Boltzmann-Gibbs distributions because of the microcanonical response matrix χ^{ij}_s turns a non positive defined matrix.

Claim 5: *The unstable region of the Boltzmann-Gibbs distributions* $\mathcal{A}^{BG}$ *corresponds to macrostates with anomalous values in the response functions, such as macrostates with negative heat capacities at constant magnetic* $C_p < 0$ *or negative isothermal compressibilities* $K_T < 0$ *in a fluid system, or macrostates with negative heat capacities at constant magnetic field* $C_H < 0$ *or isothermal magnetic susceptibilities* $\chi_T < 0$ *in a paramagnetic system.*

As already shown, the presence of macrostates with anomalous values of response functions is incompatible with the imposition of the restriction $\delta\beta^\omega_i = 0$ into the fluctuation relation (48). However, this same fluctuation relation can be rewritten to obtain an extension of Eq.(64), which is expressed here in the following:

Theorem: *The system fluctuating behavior* $C^{ij} = \left\langle \delta I^k \delta I^j\right\rangle$ *is determined within the Gaussian approximation by its response* $\chi^{ij}_\omega = -\partial\left\langle I^j\right\rangle/\partial\left\langle\beta^\omega_i\right\rangle$ *and the incidence of correlative effects* $M^j_i = \langle\delta\beta^\omega_k \delta I^j\rangle$ *that appear as a consequence of the external influence of the environment through the following fluctuation relations:*

$$\chi^{ij}_\omega = C^{ij} + \chi^{ik}_\omega M^j_k. \tag{69}$$

Eq.(69) constitutes the central result of this work. Since the fluctuation relation (62) is simply the equilibrium version of the so-called *fluctuation-dissipation theorem*, the expression (69) can be unambiguously referred to as the *extended equilibrium fluctuation-dissipation theorem*. The same one accounts for the system fluctuating behavior beyond the equilibrium thermodynamic conditions associated with the Boltzmann-Gibbs distributions (5) in a way that it is compatible with the existence of anomalous response functions. As already discussed, its validity relies on the applicability of the Gaussian approximation, which arises as a licit treatment in the limit of large system size N. As expected, the present theorem comprises all those preliminary conclusions obtained along this work, with the exception of claim 1, which was obtained from a more general equilibrium fluctuation relation, Eq.(35).

Let us finally obtain the form of the extended equilibrium fluctuation-dissipation theorem (69) in terms of the ordinary thermodynamic variables and potentials of statistical mechanics and thermodynamics: (U,β) and (X,Y) and the Enthalpy $G = U + YX$. The known relations:

$$\begin{gathered}
-\partial_{\beta}\log Z\left(\beta,Y\right)=\left\langle G\right\rangle,\ \beta\left\langle X\right\rangle=-\partial_{Y}\log Z\left(\beta,Y\right)\\
\partial_{\beta}^{2}\log Z\left(\beta,Y\right)=\left\langle \delta Q^{2}\right\rangle,\ \partial_{Y}^{2}\log Z\left(\beta,Y\right)=\beta^{2}\left\langle \delta X^{2}\right\rangle,\\
\partial_{\beta}\partial_{Y}\log Z\left(\beta,Y\right)=\beta\left\langle \delta Q\delta X\right\rangle,
\end{gathered} \tag{70}$$

allow to re-express Eq.(62) in a compact matrix form as follows:

$$\mathcal{R}=\mathcal{F}, \tag{71}$$

where the response and the correlation matrixes $\mathcal{R}$ and $\mathcal{F}$ are given by:

$$\begin{aligned}
\mathcal{R}&=-\left(\begin{array}{cc}
\partial_{\beta}\left\langle G\right\rangle & \partial_{\beta}\left(\beta\left\langle X\right\rangle\right)\\
\partial_{Y}\left\langle G\right\rangle & \beta\partial_{Y}\left\langle X\right\rangle
\end{array}\right),\\
\mathcal{F}&=\left(\begin{array}{cc}
\left\langle \delta Q^{2}\right\rangle & \beta\left\langle \delta Q\delta X\right\rangle\\
\beta\left\langle \delta X\delta Q\right\rangle & \beta^{2}\left\langle \delta X^{2}\right\rangle
\end{array}\right).
\end{aligned} \tag{72}$$

It is important to keep in mind that the Enthalpy fluctuation δG within the Boltzmann-Gibbs distribution (5) is amount of heat δQ exchanged between the system and its environment at the equilibrium, $\delta G = \delta U + Y\delta X = \delta Q$. Such a relationship does not hold in general, since $\delta G = \delta Q + X\delta Y$ when the fluctuations δY are taken into consideration. The matrix form of the fluctuation-dissipation relations (62) in terms of the conjugated thermodynamic variables (U,β) and $(X;\xi)$ with $\xi = \beta Y$ reads as follows:

$$\chi=C, \tag{73}$$

where the response and the correlation matrixes χ and C are given by:

$$\chi=-\left(\begin{array}{cc}
\partial_{\beta}\left\langle U\right\rangle & \partial_{\beta}\left\langle X\right\rangle\\
\partial_{\xi}\left\langle U\right\rangle & \partial_{\xi}\left\langle X\right\rangle
\end{array}\right), \tag{74}$$

$$C=\left(\begin{array}{cc}
\left\langle \delta U^{2}\right\rangle & \left\langle \delta U\delta X\right\rangle\\
\left\langle \delta X\delta U\right\rangle & \left\langle \delta X^{2}\right\rangle
\end{array}\right). \tag{75}$$

One can verify the existence of the following transformation rules between these two representations:

$$\mathcal{F}=\mathcal{T}C\mathcal{T}^{T},\ \mathcal{R}=\mathcal{T}\chi\mathcal{T}^{T}, \tag{76}$$

where the transformation matrix $\mathcal{T}$ is given by:

$$\mathcal{T}=\left(\begin{array}{cc} 1 & Y \\ 0 & \beta \end{array}\right). \tag{77}$$

Finally, one can rewrite the compact matrix form of extended equilibrium fluctuation-dissipation theorem (69) in these equivalent representations as follows:

$$\chi=C+\chi M \Leftrightarrow \mathcal{R}=\mathcal{F}+\mathcal{R}\mathcal{M} \tag{78}$$

where the correlation matrixes M and $\mathcal{M}$ are given by:

$$\begin{gathered} M=\left(\begin{array}{cc} \langle\delta\beta^{\omega}\delta U\rangle & \langle\delta\beta^{\omega}\delta X\rangle \\ \langle\delta\xi^{\omega}\delta U\rangle & \langle\delta\xi^{\omega}\delta X\rangle \end{array}\right), \\ \mathcal{M}=\left(\begin{array}{cc} \langle\delta\beta^{\omega}\delta Q\rangle & \beta\left\langle\delta\beta^{\omega}\delta X\right\rangle \\ \langle\delta Y^{\omega}\delta Q\rangle & \beta\left\langle\delta Y^{\omega}\delta X\right\rangle \end{array}\right), \end{gathered} \tag{79}$$

which are related by the transformation rule:

$$\mathcal{M}=\left(\mathcal{T}^{T}\right)^{-1} M\mathcal{T}^{T}. \tag{80}$$

As expected, the quantities $(\beta^{\omega}, Y^{\omega})$ denote the thermodynamic parameters of the environment.

3. Examples of application

3.1. Application to fluid and magnetic systems

Direct application of Eqs.(72) and (79) to the thermodynamic variables of a fluid system with Enthalpy $G=U+pV$ leads to the following expressions of the matrixes $\mathcal{R}$, $\mathcal{F}$ and $\mathcal{M}$:

$$\begin{gathered} \mathcal{R}=\left(\begin{array}{cc} T^{2}C_{p} & V\left(T\alpha_{p}-1\right) \\ -\left(\partial G/\partial p\right)_{T} & \beta V K_{T} \end{array}\right), \\ \mathcal{F}=\left(\begin{array}{cc} \langle\delta Q^{2}\rangle & \beta\left\langle\delta Q\delta V\right\rangle \\ \beta\left\langle\delta V\delta Q\right\rangle & \beta^{2}\left\langle\delta V^{2}\right\rangle \end{array}\right), \\ \mathcal{M}=\left(\begin{array}{cc} \langle\delta\beta^{\omega}\delta Q\rangle & \beta\left\langle\delta\beta^{\omega}\delta V\right\rangle \\ \langle\delta p^{\omega}\delta Q\rangle & \beta\left\langle\delta p^{\omega}\delta V\right\rangle \end{array}\right), \end{gathered} \tag{81}$$

where $C_{p}=\left(\partial G/\partial T\right)_{p}$ and $\alpha_{p}=V^{-1}\left(\partial V/\partial T\right)_{p}$ are the heat capacity and the thermal expansibility at constant pressure, while $K_{T}=-V^{-1}\left(\partial V/\partial p\right)_{T}$ is the isothermal compressibility. The symmetry of the response matrix in Eq.(81) leads to the thermodynamical identity:

$$\left(\frac{\partial G}{\partial p}\right)_{T}=V-T\left(\frac{\partial V}{\partial T}\right)_{p}. \tag{82}$$

The corresponding expressions for a magnetic system with Enthalpy $G = U - HM$ are directly derived from the previous results by substituting $(V, p) \to (M, -H)$:

$$\mathcal{R} = \left(\begin{array}{cc} T^2 C_H & T\left(\partial M/\partial T\right)_H - M \\ \left(\partial G/\partial H\right)_T & \beta\chi_T \end{array} \right),$$
$$\mathcal{F} = \left(\begin{array}{cc} \langle \delta Q^2 \rangle & \beta \langle \delta Q \delta M \rangle \\ \beta \langle \delta M \delta Q \rangle & \beta^2 \langle \delta M^2 \rangle \end{array} \right), \qquad (83)$$
$$\mathcal{M} = \left(\begin{array}{cc} \langle \delta\beta^{\omega} \delta Q \rangle & \beta \langle \delta\beta^{\omega} \delta M \rangle \\ -\langle \delta H^{\omega} \delta Q \rangle & -\beta \langle \delta H^{\omega} \delta M \rangle \end{array} \right),$$

where $C_H = \left(\partial G/\partial T\right)_H$ is the heat capacity at constant magnetic field and $\chi_T = \left(\partial M/\partial H\right)_T$ the isothermal magnetic susceptibility. The symmetry of the response matrix in Eq.(83) leads to the thermodynamical identity:

$$\left(\frac{\partial G}{\partial H}\right)_T = T\left(\frac{\partial M}{\partial T}\right)_H - M. \qquad (84)$$

3.2. Thermodynamic stability of macrostates with anomalous response functions

A direct consequence of extended equilibrium fluctuation-dissipation theorem (78) are the *conditions of thermodynamic stability*, which can be compatible in this framework with the existence of anomalous response functions. In the present subsection, let us discuss two simple application examples involving macrostates with negative heat capacities and a numerical testing during the study of macrostates with anomalous values in the isothermal susceptibility of a paramagnetic system.

3.2.1. Thirring's constraint: Let us first consider an equilibrium situation involving only the conjugated thermodynamic pair energy-temperature. Since $\delta X = 0 \Rightarrow \delta Q \equiv \delta U$, Eq.(78) drops to the generalized equilibrium fluctuation-dissipation relation involving the heat capacity (3) already obtained in our previous work [10]. A simple way to implement the existence of non-vanishing correlated fluctuations $\langle \delta\beta^{\omega} \delta U \rangle$ is by considering a thermal contact with a bath (B) having a finite heat capacity $C_B > 0$. Clearly, the bath experiences a temperature fluctuation δT_B when the system absorbs or releases an amount of energy δU, $\delta T_B = -\delta U/C_B \Rightarrow \langle \delta\beta_B \delta U \rangle \neq 0$. Eq.(3) can be rephrased as follows:

$$\frac{C C_B}{C + C_B} = \beta^2 \left\langle \delta U^2 \right\rangle \qquad (85)$$

after considering the Gaussian approximation $\delta\beta_B = \beta^2 \delta U / C_B$ (here $k_B \equiv 1$). Since the expectation value of the square dispersion of the energy fluctuations $\langle \delta U^2 \rangle$ is nonnegative, the thermodynamic stability of macrostates with negative heat capacities $C < 0$ demands the applicability of the following condition:

$$\frac{C C_B}{C + C_B} > 0 \Rightarrow 0 < C_B < |C|. \qquad (86)$$

This result is same constraint derived by Thirring in the past [39]. As already commented, the existence of anomalous macrostates with negative heat capacities is a distinguish feature of many nonextensive systems. However, since the derivation of Thirring's constraint (86) demands the existence of a thermal contact, such a stability condition could be only applicable to short-range interacting systems whose sizes are large enough in order to dismiss the surface energy involved in their thermodynamic interaction.

3.2.2. Stability condition in an astrophysical situation: Let us now consider a paradigmatic situation with long-range interactions: the astrophysical system composed by a globular cluster (C) and its nearby galaxy (G). Let us denote by C_C and C_G their respective heat capacities, being $|C_C| \ll |C_G|$. At first glance, the galaxy could be taken as a bath with a very large heat capacity, which naively suggests that the thermo-statistical description of the globular cluster could be performed with the help of the Gibbs' canonical ensemble (2). In this context, however, the known exigence of a *thermal contact* fails as a consequence of the long-range nature of Gravitation.

Astronomers known since the late 1800s that the attractive character and power-law $1/r^2$ dependence of gravitational interaction favor a concentration of stars towards the inner regions of any astrophysical structure and the increasing of their kinetic energies while a decreasing of the total energy [40], a tendency stimulating the development of distribution profiles with a *core-halo structure* and negative heat capacities. Consequently, it is a rule rather than an exception the fact that the globular cluster and its nearby galaxy simultaneously exhibit macrostates with negative heat capacities, $C_C < 0$ and $C_G < 0$. Since Thirring's constraint (86) cannot be satisfied, and overall, the same one is inapplicable in the present situation, the energy associated with the thermodynamic interaction between the globular cluster and the galaxy should play a fundamental role in the thermodynamic stability.

To obtain a general idea about how works the above mechanism, let us denote by W_{G-C} the energy contribution of the total energy $U_T = U_G + U_C + W_{G-C}$ accounting for the potential energy V_{G-C} associated with the gravitational interaction between these systems, as well as the kinetic energy contributions of the collective translational motion T_{G-C}:

$$W_{G-C} = T_{G-C} + V_{G-C}. \tag{87}$$

In general, W_{G-C} should be significantly smaller than the internal energy of the galaxy U_G, but it could be comparable to the internal energy of the globular cluster U_C, $|W_{G-C}| \sim |U_C| \ll |U_G|$. Strictly speaking, the total energy U_T changes as a consequence of gravitational influence of other neighboring astrophysical structures and the evaporation of stars. However, the composed system globular cluster-galaxy can arrive at a certain *quasi-stationary evolution* under its own relaxation mechanisms (collisions, violent relaxation, etc. [33]) and the incidence of stars evaporation, where U_T changes at a very slow rate, $\dot{U}_T \simeq 0$. In these conditions, it is possible to speak about the existence of a certain *metastable thermodynamic equilibrium* between the globular cluster and its nearby galaxy, a fact that it is manifested in the observation of universal profiles such as the Vaucouleurs' law in the surface brightness of the elliptical galaxies and the Michie-King profiles for the distribution of stars in the globular clusters [41].

Due to the coupling between collective and internal degrees of freedom, the energy contribution W_{G-C} will depends on the internal states of the globular cluster and its nearby galaxy, so that, it can be expressed by their respective internal energies (U_G, U_C) as well as some other collective variables σ (their total masses, relative separation, etc.), $W_{G-C} \simeq W(U_G, U_C; \sigma)$. Consequently, the cluster and galaxy undergo, in general, a *nonlinear energetic interchange* during their thermodynamic interaction, $\Delta U_G \neq -\Delta U_C$, since part of the energy transferred from the galaxy ΔU_G towards the cluster is used to modify the interaction energy W_{G-C}. Such a nonlinear interchange contracts with the usual thermal contact between a Gibbs' bath (B) and an extensive system (S), where $\Delta U_B = -\Delta U_S$.

As a consequence of the separability as well as the small number of degrees of freedom of collective motions $N_{G-C}^{collective}$ in comparison with the internal degrees of freedom of the galaxy $N_G^{internal}$ and the globular cluster $N_G^{internal}$, the distribution associated with the present situation can be approximately given as the product of their respective densities of states Ω_G and Ω_C:

$$\rho(U_C) \simeq A\Omega_G(U_G)\Omega_C(U_C), \tag{88}$$

where the internal energies U_G and U_C obey the constraint:

$$U_T = U_G + U_C + W_{G-C} \sim const, \tag{89}$$

with A being a normalization constant. Since the probabilistic weight $\omega(U_C) \sim \Omega_G(U_G)$, the environment inverse temperature (see definition in Eq.(15)) is given by:

$$\beta^{\omega} = -\left.\frac{\partial S_G(U_G)}{\partial U_C}\right|_{U_T} = \beta_G \alpha, \tag{90}$$

where the subindex U_T means that the partial derivation is taken at constant total energy. In this last expression, $\beta_G = \partial S_G(U_G)/\partial U_G$ represents the ordinary microcanonical inverse temperature of the galaxy and $\alpha = \alpha(U_C)$:

$$\alpha = -\left.\frac{\partial U_G}{\partial U_C}\right|_{U_T} \equiv 1 + \left.\frac{\partial W_{G-C}}{\partial U_C}\right|_{U_T} = \alpha(U_C), \tag{91}$$

a dimensionless factor that accounts for the underlying nonlinear energetic interchange. Clearly, the inverse temperature (90) is an *effective concept* that characterizes how the globular cluster *fills* the thermodynamic influence of the galaxy. Since in general $\alpha \neq 1$, the condition of thermal equilibrium $\beta^{\omega} = \beta_C$ implies that the microcanonical inverse temperatures of the galaxy and the globular cluster are different, $\beta_G \neq \beta_C$.

Since the heat capacity of galaxy C_G is extremely large in comparison with the heat capacity of the globular cluster C_C, the inverse temperature β_G can be taken as a constant parameter. Thus, we can limit to consider the thermal fluctuations of effective inverse temperature β^{ω} associated with the nonlinear energetic interchange:

$$\delta\beta^{\omega} \simeq \beta_G \alpha'(U_C)\delta U_C. \tag{92}$$

By substituting this last approximation into fluctuation-dissipation relation of Eq.(3) and considering the condition of thermal equilibrium $\beta^{\omega} = \beta_C$, one obtains:

$$\frac{C_C}{1 + (\alpha\beta_C)^{-1} C_C \alpha'} = \beta_C^2 \left\langle \delta U_C^2 \right\rangle. \tag{93}$$

Since $\langle \delta U_C^2 \rangle > 0$, one arrives at the following condition of thermodynamic stability:

$$\beta_C C_C^{-1} + \alpha^{-1}\alpha' > 0. \tag{94}$$

According to the present result, the globular cluster with $C_C < 0$ can be found in a metastable thermodynamic with its nearby galaxy provided that the underlying nonlinear energetic interchange associated with the interaction energy W_{G-C} ensured the applicability of stability condition (94).

3.2.3. Anomalous response in paramagnetic systems: Besides the long-range interacting systems as the astrophysical example previously discussed, the existence of macrostates with negative heat capacities is a relevant feature indicating the occurrence of a *discontinuous* (first-order) *phase transition* in a mesoscopic short-range interacting system [8]. The observation of anomalies in discontinuous phase transitions is not restricted to the heat capacity, but other response functions can also exhibit anomalous values too [32]. In fact, their presence could be the origin of the sudden change experienced by a given physical observable X with a small varying of its conjugated control parameter Y below the critical temperature T_c.

A typical example is found in a magnetic system between the magnetization M and the external magnetic field H. The sudden inversion of magnetization $M \mapsto -M$ with $M \neq 0$, observed in turns to the value $H = 0$ below the critical temperature T_c, is a consequence of the existence of unstable diamagnetic states $\chi_T < 0$. Such states are anomalous for magnetic materials undergoing a paramagnetic-ferromagnetic phase transition, in those equilibrium situations where the fluctuation-dissipation relation $\chi_T = \beta \langle \delta M^2 \rangle$ is supposed to be relevant, and the inequality $\chi_T > 0$ arises as the ordinary stability condition.

Let us consider again a magnetic system with internal energy U and magnetization M. As already shown in this work, the corresponding form of the generalized equilibrium fluctuation-dissipation relations $\mathcal{R} = \mathcal{F} + \mathcal{R}\mathcal{M}$ is specified by matrixes of Eq.(83). In the sake of simplicity, let us restrict to a particular equilibrium situation where temperature of environment β^{ω} shows a constant value β, but the external magnetic field H^{ω} acting on the magnetic system undergoes non-vanishing thermal fluctuations $\langle \delta H^{\omega} \delta M \rangle \neq 0$.

The simplest way to implement the above behavior is by admitting the existence small thermodynamic fluctuation of the magnetic field H^{ω} in turn to a constant value H coupled to the total magnetization of the system:

$$H^{\omega} = H - \lambda \delta M / N, \tag{95}$$

where N is the system size and λ a suitable coupling constant. The general equilibrium fluctuation-dissipation relation involving the isothermal magnetic susceptibility χ_T :

$$\beta\chi_T = \beta^2\left\langle\delta M^2\right\rangle + \left(T\left(\partial M/\partial T\right)_H - M\right)\beta\left\langle\delta\beta^\omega\delta M\right\rangle - \beta^2\chi_T\left\langle\delta H^\omega\delta M\right\rangle \tag{96}$$

drops to:

$$\beta\chi_T = \beta^2\left\langle\delta M^2\right\rangle - \beta^2\chi_T\left\langle\delta H^\omega\delta M\right\rangle \tag{97}$$

in the present equilibrium situation, which can be rewritten as follows:

$$\beta\left\langle\delta M^2\right\rangle = \frac{\chi_T}{1+\lambda\chi_T/N}. \tag{98}$$

Since the standard deviation of magnetization is nonnegative, $\left\langle\delta M^2\right\rangle > 0$, the anomalous macrostates with negative values of isothermal magnetic susceptibility $\chi_T < 0$ are thermodynamically stable when the condition:

$$\lambda + N/\chi_T > 0 \tag{99}$$

holds. Eq.(98) can also be used to obtain the dispersion of the external magnetic field H^ω:

$$\left\langle\left(\delta H^\omega\right)^2\right\rangle = \frac{1}{N^2}\lambda^2\left\langle\delta M^2\right\rangle = \frac{1}{N\beta}\lambda^2\frac{\chi_T/N}{1+\lambda\chi_T/N}. \tag{100}$$

Since χ_T grows with N in general as $\chi_T \propto N$, the dispersions of the magnetization and the external magnetic field behave as $\left\langle\delta M^2\right\rangle \propto N$ and $\left\langle\left(\delta H^\omega\right)^2\right\rangle \propto 1/N$. When the system size N is large enough, the fluctuations of the external magnetic field H^ω are very small. From the viewpoint of thermodynamics (where thermal fluctuations are dismissed due to the large size of conventional thermodynamic systems), the present equilibrium does not differ from conventional situations where the external magnetic field is assumed to be constant. However, it is easy to see that both the fluctuating behavior accounted for expressions (98) and (100), as well as stability condition (99) crucially depend on the external conditions which have been imposed to, which are specified here by the coupling constant λ.

According to the stability condition (99), macrostates with negative isothermal susceptibilities can be found in a stable equilibrium with an appropriate choosing of the coupling constant λ. Even, fluctuation relation (98) could be also used to obtain the isothermal magnetic susceptibility per particle $\bar{\chi}_T = \chi_T/N$ in terms of fluctuating behavior of magnetization:

$$\bar{\chi}_T^{-1} = \frac{1+\beta\left\langle\delta H^\omega\delta M\right\rangle}{\beta\left\langle\delta M^2\right\rangle}N \equiv \frac{1-\lambda\beta\sigma_m^2}{\beta\sigma_m^2}, \tag{101}$$

where $\sigma_m^2 = \left\langle\delta M^2\right\rangle/N$ represents the thermal dispersion of magnetization.

Certainly, we unknown how the external magnetic influence expressed in Eq.(95) could be implemented in a real experimental situation. Fortunately, their consequences are very easy to check with the help of Monte Carlo simulations. Let us consider the known 2D Ising model on the square lattice $L\times L$ with periodic boundary condition:

$$U = -\sum_{\langle ij\rangle} s_i s_j,\ M = \sum_i s_i, \tag{102}$$



Figure 1. Isotherms of 2D Ising model with $L = 25$ derived from Metropolis Monte Carlo simulations by using the acceptance probability (103): Panel a) magnetic field $\langle H^{\omega}\rangle$ *versus* magnetization $\langle m\rangle$; Panel b) inverse isothermal magnetic susceptibility $\bar{\chi}_T^{-1}$ *versus* magnetization $\langle m\rangle$. The labels in both panels indicate the value of β for the corresponding isotherm.

where the sum $\langle ij\rangle$ considers nearest-neighbor interactions only, and the spin variables $s_i = \pm 1$. The above equilibrium conditions can be easily implemented by using a Metropolis method with acceptance probability $p = p\left(U, M\left|U + \Delta U, M + \Delta M\right.\right)$:

$$p = \min\left\{\exp\left[-\beta\Delta U + \beta H^{\omega}\Delta M\right]\right\}, \tag{103}$$

which is a particular expression of Eq.(18). By denoting $m = M/N$ the magnetization per particle, the external magnetic field in this study is given by $H^{\omega} = \bar{H} + \lambda\left(m - \bar{m}\right)$, where $\bar{m}$ and $\bar{H}$ are suitable estimation of the expectation values $\langle m\rangle$ and $\langle H^{\omega}\rangle$. Our interest is to obtain the isotherms of 2D Ising model within anomalous regions with $\chi_T < 0$. Thus, the values of parameters $\left(\bar{H}, \bar{m}\right)$ can iteratively provide by using the isothermal magnetic susceptibility per particle $\bar{\chi}_T$ of a previous simulation throughout the expression:

$$\bar{H}_{i+1} = \bar{H}_i + \left(\bar{\chi}_T\right)_i^{-1}\left(\bar{m}_{i+1} - \bar{m}_i\right), \tag{104}$$



Figure 2. Isotherms for $L = 25$ and $\beta = 0.46$ obtained from Metropolis Monte Carlo simulations based on the Boltzmann-Gibbs distributions (BG) where $H^{\omega} = H \sim const \rightarrow \langle \delta H^{\omega} \delta M \rangle = 0$ and the present Metropolis Monte Carlo simulations with $\langle \delta H^{\omega} \delta M \rangle \neq 0$.

where the step $\Delta m = \bar{m}_{i+1} - \bar{m}_i$ should be small. The initial value $\bar{m}_0$ is assumed to be the average of magnetization obtained from an ordinary Metropolis Monte Carlo simulation with $H^{\omega} = \bar{H}_0 \sim const$ ($\lambda \equiv 0$) far enough from the unstable region with $\chi_T < 0$, $\bar{m}_0 = \langle m \rangle$.

Although any real value of coupling constant λ satisfying stability condition (99) is admissible, it is desirable to impose some criterium in order to reduce thermal dispersions of the system magnetization (98) and the external magnetic field (100). Our interest is to obtain the isotherms in the $H - M$ diagram, and hence, an optimal value for λ can be chosen in order to minimize the total dispersion $\sigma^2 = \sigma_H^2 + \sigma_m^2$, where:

$$\sigma_H^2 = N \left\langle (\delta H^{\omega})^2 \right\rangle \tag{105}$$

is the thermal dispersion of the external magnetic field. The analysis yields:

$$\lambda_{op} = \sqrt{a^2 + 1} - a, \tag{106}$$

where a is the inverse of the isothermal magnetic susceptibility per particle, $a = \bar{\chi}_T^{-1}$, whose value can be estimated from the previous simulation as the case of parameters $\left(\bar{H}, \bar{m}\right)$ in Eq.(104). Notice that this optimal value guarantees the validity of stability condition (99).

Results of extensive Monte Carlo simulations by using the procedure explained above are shown in FIG.1. We have used a square lattice with $L = 25$ and considered $n = 10^6$ iterations for each calculated point. This results clearly revealed the presence of anomalous diamagnetic states $\bar{\chi}_T < 0$ for inverse temperatures β above the critical point $\beta_c^{L=25} \simeq 0.41$. We only show here macrostates with $\langle m \rangle > 0$ due to the underlying symmetry $M \rightarrow -M$ and $H \rightarrow -H$ of the Enthalpy $G = U - HM$.

As clearly shown in FIG.2, while the 2D Ising model isotherms are able to describe a backbending in the $H - M$ diagram under a magnetic influence obeying stability condition (99), they merely undergo a sudden jump in magnetization in turns to the point $H = 0$ under the influence of a constant magnetic field, that is, a situation with $\lambda = 0$. As already commented, such a sudden jump is interpreted as the occurrence of a discontinuous phase transition in conventional statistical mechanics, whose existence is a direct manifestation of the presence of an unstable branch with negative susceptibilities $\chi_T < 0$. Our Monte Carlo simulations show that such an anomalous region can turn accessible with an appropriate selection of the external experimental conditions. The present results can be extended to the case of fluid systems. In fact, the behavior of the 2D Ising model isotherms is fully analogous to the well-known backbending of Van der Waal gas isotherms in the $p - V$ diagram, which is associated with macrostates with negative isothermal compressibilities $K_T < 0$ [42].

4. Final remarks

We have obtained in this work a generalization of equilibrium fluctuation-dissipation relations compatible with the existence of macrostates with anomalous response functions, Eq.(69). Such a development has been carried out by starting from a generic extension of the Boltzmann-Gibbs distribution and its control thermodynamic parameters (β, Y) that characterize the external thermodynamic influence of an environment, Eqs.(7) and (15). Since the environment thermodynamic control parameters turn, in general, *fluctuating quantities*, a new operational definition (66) for the concept of response function has been also considered. Remarkably, the physics behind the generalized equilibrium fluctuation-dissipation relations (69) is quite simple. The key ingredient relies on the that the environment thermodynamic parameters $\{\beta_i^e\}$ undergo in general correlated fluctuations with the system observables $\{I^i\}$ as a consequence of the underlying thermodynamic interaction. Such a realistic behavior, systematically disregarded by the using of Boltzmann-Gibbs distributions (6), leads to the consideration of a new correlation matrix $\mathcal{M}$ in the equilibrium fluctuation-dissipation relations, Eq.(79), which characterizes the existence of these correlative effects. The new theorem explicitly expresses that the system fluctuating behavior and its thermodynamic stability crucially depend on the nature of the external influence of the environment.

As example of applications, we have obtained particular expressions of the generalized equilibrium fluctuation-dissipation relations for a fluid and magnetic systems. After, we have derived two criteria of thermodynamic stability accounting for the existence of negative heat capacities: Eq.(86), for systems with short-range interactions ensuring the existence of a *thermal contact*; and Eq.(94), for a metastable equilibrium involving long-range interactions between a globular cluster and its nearby galaxy. Finally, we have arrived at a stability condition compatible with anomalous isothermal susceptibilities associated with the occurrence of a discontinuous phase transition in magnetic systems, Eq.(99), whose consequences have been also numerically tested throughout Monte Carlo simulations.

We would like mention some open questions deserving a special attention in future works:

- The usual equilibrium fluctuation-dissipation relations (71) constitute a particular expression of the so-called *fluctuation-dissipation theorem*:
$$\beta C^{ij}\left(\tau\right)=\frac{1}{i\pi}P\int_{-\infty}^{+\infty}\frac{\chi^{ij}\left(\omega\right)}{\omega}\cos\left(\omega\tau\right)d\omega, \tag{107}$$
relating correlations and response functions in linear response theory [42]. The present generalization of equilibrium fluctuation-dissipation relations inspired a corresponding extension of this last fundamental result.
- The numerical study of the 2D Ising model clearly shows that the present ideas are very useful in order to implement some new Monte Carlo algorithms inspired on statistical mechanics [12] in order to access to macrostates with anomalous response functions.
- At first glance, rigorous fluctuations relations (35) and (36) clearly exhibit a *tensorial character*, which suggests a direct connection of the present approach with some geometric formulations of fluctuation theory developed in the past [22, 23]. A preliminary analysis of such a question is presented in a recent paper [11].

Acknowledgements

It is a pleasure to acknowledge partial financial support by FONDECYT 3080003. Velazquez also thanks the partial financial support by the project PNCB-16/2004 of the Cuban National Programme of Basic Sciences.

References

[1] Lynden-Bell D 1967 *Mon. Not. R. Astro Soc.* **136**, 101; Lynden-Bell D and Wood R 1968 *Mon. Not. R. Astro Soc.* **138** 495.
[2] Moretto L G, Ghetti R, Phair L, Tso K and Wozniak, G.J. 1997 *Phys. Rep.* **287** 250.
[3] D'Agostino *et al* 2000 *Phys. Lett.* **B 473** 219.
[4] Gross D H E and Madjet M E 1997 *Z. Phys.* **B 104** 521.
[5] Lynden-Bell D and Lynden-Bell R M 1977 *Mon. Not. R. Astro Soc.* **181** 405.
[6] Padmanabhan T 1990 *Physics Reports* **188** 285.
[7] Lynden-Bell D 1999 *Physica A* **26** 293.
[8] Gross D H E 2001 *Microcanonical thermodynamics: Phase transitions in Small systems*, *66 Lectures Notes in Physics* (World scientific, Singapore).
[9] Dauxois T, Ruffo S, Arimondo E and Wilkens M (Eds.) 2002 *Dynamics and Thermodynamics of Systems with Long Range Interactions*, *Lecture Notes in Physics* (Springer, New York).
[10] Velazquez L and Curilef S 2009 *J. Phys. A: Math. Theor.* **42** 095006.
[11] Velazquez L and Curilef S 2009 *J. Phys. A: Math. Theor.* **42** 335003.
[12] Landau P D and Binder K 2000 *A guide to Monte Carlo simulations in Statistical Physics* (Cambridge Univ. Press).
[13] Bohr N in: *Collected Works*, Kalckar J (Ed) 1985 (North-Holland, Amsterdam), Vol. 6, pp. 316-330, 376-377.
[14] Heisenberg W 1969 *Der Teil und das Gauze*, Ch. 9. R. piper, Miinchen.

[15] Rosenfeld L in: *Ergodic Theories*, P. Caldirola (Ed) 1961 (Academic Press, New York) pp. 1.

[16] Mandelbrot B 1956 *IRE Trans . Inform. Theory* **IT-2** 190.

[17] Gilmore R 1985 *Phys. Rev. A* **31** 3237.

[18] Lindhard J in: *The Lessons of Quantum Theory* de Boer J, Dal E and Ulfbeck O (Ed) 1986 (Elsevier, Amsterdam).

[19] Lavenda B 1987 *Int. J. Theor. Phys.* **26** 1069; 1991 *Statistical Physics: A Probabilistic Approach* (Wiley, New York).

[20] Schölg F 1988 *J. Phys. Chem. Sol.* **49** 679.

[21] Uffink J and van Lith J 1999 *Found. Phys.* **29** 655.

[22] Weinhold F 1975 *J. Chem. Phys. Vol* **63** 22479; 2484; 2488; 2496.

[23] Ruppeiner G 1995 *Rev. Mod. Phys.* **67** 605 and references therein.

[24] Landau L D and Lifzhitz E M 1977 *Statistical Physics* (Pergamon, New York).

[25] Antoni M and Ruffo S 1995 *Phys. Rev. E* **52** 2361.

[26] Latora V, Rapisarda A and Ruffo S 1999 *Physica* **D 131** 38; 2000 *Prog. Theor. Phys. Suppl.* **139** 204; 2001 *Nucl. Phys. A* **681** 331c.

[27] Ison M J, Chernomoretz A and Dorso C O 2004 *Physica* **A 341** 389.

[28] Einarsson B 2004 *Phys. Lett.* **A 332** 335.

[29] Chomaz P and Gulminelli F 2006 *Eur. Phys. J.* **A 30** 317.

[30] Gulminelli F 2007 *Nuclear Physics* **A 791** 165.

[31] Lovett R 2007 *Rep. Prog. Phys.* **70** 195.

[32] Campa A, Ruffo S and Touchette S 2007 *Physica* **A 385** 233.

[33] Chavanis P H in: *Dynamics and Thermodynamics of Systems with Long Range Interactions*, Lecture Notes in Physics, Dauxois T, Ruffo S, Arimondo E and Wilkens M (Ed) 2002 (Springer, New York); e-print (2002) [cond-mat/0212223].

[34] Beck C 2001 *Phys. Rev. Lett.* **87** 18061; 2000 *Physica* **A 277** 115; 2000 *Physica* **A 286** 164.

[35] Cugliandolo L F 2002 *J. Non-Cryst. Solids* **307-310** 161-171.

[36] Beck C and Cohen E G D 2003, *Physica* **A 322** 267; Cohen E G D 2004, *Physica* **D 193** 35.

[37] Metropolis N, Rosenbluth A W, Rosenbluth M N, Teller A H and Teller E 1953 *J. Chem. Phys.* **21** 1087.

[38] Velazquez L and Curilef S 2009 *Uncertainty relations of Statistical Mechanics* To appear in *Modern Physics Letters B.*

[39] Thirring W 1980 *Quantum Mechanics of large systems* (Springer) Ch. 2.3.

[40] Eddington A S 1926 *The Internal Constitution of the Stars*, (Cambridge); 1916 *Mon. Not. R. Astr. Soc.* **76** 525.

[41] Binney J and Tremaine S 1987 *Galactic Dynamics*, Princeton Series in Astrophysics, (Princeton, NJ).

[42] Reichl L E 1980 *A modern course in Statistical Mechanics*, (Univ. Texas Press, Austin).